

IMPACT PICTURE AND THE RATIO OF ELASTIC
TO TOTAL CROSS SECTION AT ISR ENERGIES

H. Cheng

Massachusetts Institute of Technology, Cambridge, Massachusetts

J. K. Walker

National Accelerator Laboratory, Batavia, Illinois

and

T. T. Wu

Harvard University, Cambridge, Massachusetts

March 1974



IMPACT PICTURE AND THE RATIO OF ELASTIC
TO TOTAL CROSS SECTION AT ISR ENERGIES

Hung Cheng^{*}

Department of Mathematics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139

James K. Walker[†]

Physics Department, National Accelerator Laboratory,
Batavia, Illinois 60510

Tai Tsun Wu[‡]

Gordon McKay Laboratory, Harvard University,
Cambridge, Massachusetts 02138

Abstract

We show that the impact picture of the pomeron together with f^0 and ω exchange account for the shrinkage anti-shrinkage of the elastic forward peak for proton-proton and antiproton-proton scattering for energies greater than 20 GeV. We predict that the ratio of the elastic to total pp cross section rises by 7% in the energy range available at the CERN ISR.

In 1970, we described the implications of relativistic quantum field theory for scattering at very high energy. We called these general qualitative features the Impact Picture of high energy scattering.¹ Subsequently in 1972, we² made quantitative predictions of the rise in the proton-proton total cross section that would be observed at the CERN Intersecting Storage Rings. At that time we predicted

- (i) the proton-proton total cross section would increase by about 3 mb,
- (ii) a dip would develop in the elastic proton-proton differential cross section at $|t| \approx 1.3$,
- (iii) considering only the pomeron contribution, the ratio of elastic to total cross section would increase by a few percent and approach its limiting value of 0.5 only at $s \gg 10^{10}$ GeV.

Shortly thereafter, two^{3,4} independent experiments measured the total cross section and their results indicated that the proton-proton total cross section had increased by about 4 mb. In addition, experimental evidence⁵ was obtained on the existence of a deep dip in the elastic differential cross section at the predicted region of $|t|$. So far, no data, with the required accuracy, has been obtained pertaining to the third prediction. These experimental results gave considerable encouragement to the interpretation of high energy strong interactions in terms of the impact picture.

Based on the observed increase in the cross section we then made a series of predictions on the energy dependence of other hadron-hadron total

cross sections, elastic scattering,⁶ the ratio of real to imaginary parts of forward elastic amplitudes,⁷ and diffraction dissociation.⁸ Many of these predictions will shortly be confronted with experimental results from the National Accelerator Laboratory, USA.

In this paper, we reconsider in more detail the ratio of elastic to total cross section. After taking due account of Regge background terms in the elastic amplitude, thereby obtaining a more reliable estimate of the energy dependence of the elastic cross section at low energies, we predict that the ratio of elastic to total proton-proton cross section will increase by 7% in the energy range available at the ISR, namely $500 \lesssim s \text{ (GeV)}^2 \lesssim 3800$. Sufficiently accurate measurements appear feasible and constitute a critical test of the Impact Picture.

At the energies for which this prediction is made and for the accuracy which is required, our previous formulation of the model of the Impact Picture must be supplemented by the addition of some Regge concepts. We have incorporated the dominant Regge background in terms of f^0 and ω exchange. Inclusion of these exchanges in a phenomenological manner together with our previous model of the Impact Picture of the pomeron allows us to describe

1. rapid shrinkage of the pp diffraction peak at Serpukhov energies ($s \lesssim 100 \text{ GeV}^2$),
2. slow shrinkage of the pp diffraction peak at NAL and ISR energies ($100 \lesssim s \text{ (GeV)}^2 \lesssim 3000$),
3. slow anti-shrinkage of the antiproton-proton diffraction peak at Serpukhov energies ($s \lesssim 100 \text{ GeV}^2$),

4. the ratio of real to imaginary parts of the proton-proton elastic amplitude as a function of energy and momentum transfer, t .

We start by assuming that elastic hadron-hadron scattering amplitudes are dominated by three terms, namely

1. the pomeron as described and parameterized by us previously;⁶
2. tensor exchanges (f^0, A_2) with even signature and trajectory $\alpha(t) = 0.5 + t$, t in $(\text{GeV}/c)^2$;
3. vector exchanges (ρ, ω) with odd signature and trajectory degenerate with $\alpha(t)$.

For simplicity, we neglect spin-flip amplitudes, whose contributions to elastic scattering are known to be small near the forward direction. In addition we neglect the isospin exchange contributions which are known to be small from measurements on charge exchange scattering. We express the pomeron contribution to the amplitude as usual by

$$A_p(s, t) = \frac{is}{2\pi} \int dx_{\perp} e^{-i \vec{\Delta} \cdot \vec{x}_{\perp}} D(s, x_{\perp})$$

where $\vec{\Delta}$ is the momentum transfer, $-t = \Delta^2$, x_{\perp} is the impact parameter and

$$D(s, x_{\perp}) = 1 - \exp[-S(s) F(x_{\perp}^2)].$$

On the basis of our experience with field theory we have taken

$$S(s) = [E \exp(-i\pi/2)]^c / [\ln(E \exp(-i\pi/2))]^{c'}$$

where E is the incident laboratory energy in GeV, c is a real constant,

and c' has been chosen as before to be zero or one. As the predictions we obtain are similar in the two cases, we present the $c' = 0$ solution.

In addition, we have chosen

$$F(x_{\perp}^2) = f \exp [-\lambda (x_{\perp}^2 + x_0^2)^{1/2}]$$

where f , λ , and x_0 are parameters which have already been determined.⁶

The f^0 and ω type exchanges are assumed in the usual way to give rise to an amplitude of the form

$$A_{f^0}(s, t) = \gamma_f(t) \left[\frac{\alpha(t)}{\alpha(0)} \right] \left\{ \frac{1 + \exp [-i\pi \alpha(t)]}{\sin \pi \alpha(t)} \right\} s^{[\alpha(t) - 1]}$$

and

$$A_{\omega}(s, t) = \gamma_{\omega}(t) \left\{ \frac{1 - \exp [-i\pi \alpha(t)]}{\sin \pi \alpha(t)} \right\} s^{[\alpha(t) - 1]}$$

where $\gamma_{f, \omega}(t)$ is the residue function, s is in units of GeV^2 . The differential cross sections are given by

$$\frac{d\sigma}{dt} (pp) = 1.2233 \left| \frac{1}{s} A_p + A_{f^0} - A_{\omega} \right|^2 \text{ mb}/(\text{GeV}/c)^2$$

and

$$\frac{d\sigma}{dt} (\bar{p}p) = 1.2233 \left| \frac{1}{s} A_p + A_{f^0} + A_{\omega} \right|^2 \text{ mb}/(\text{GeV}/c)^2.$$

Fits by other authors⁹ to pp and $\bar{p}p$ elastic scattering at less than 20 GeV have lead to the parameterization

$$\gamma_{\omega}(t) = \bar{\gamma}_{\omega} \exp(0.5 t) (1 + t/0.14) [(1 + t)^2 + 0.16]/1.16.$$

This parameterization reproduces the well known crossover effect in the elastic differential cross section at $t = 0.14 (\text{GeV}/c)^2$ and is accurate for $|t| < 0.4 (\text{GeV}/c)^2$. We take the same residue function and apply it for energies greater than 20 GeV with no new parameters.

The residue $\gamma_f(t)$ has not previously been determined without some theoretical assumption on the pomeron contribution. For simplicity we take

$$\gamma_f(t) = -\bar{\gamma}_f \exp(dt)$$

for the small t region $0 \leq |t| \leq 0.4 (\text{GeV}/c)^2$, where d is a single adjustable parameter.

From the optical theorem we have

$$\sigma_{\text{total}}(\bar{p}p) = 4.893 \left[\frac{\text{Im } A_p(s, 0)}{s} + s^{-1/2} [\bar{\gamma}_f + \bar{\gamma}_{\omega}] \right] \text{ mb}$$

$$\sigma_{\text{total}}(pp) = 4.893 \left[\frac{\text{Im } A_p(s, 0)}{s} + s^{-1/2} [\bar{\gamma}_f - \bar{\gamma}_{\omega}] \right] \text{ mb.}$$

We have previously⁶ obtained the parameters given below for the case of nucleon-nucleon scattering:

| | |
|--|----------|
| c | 0.082925 |
| c' | 0 |
| λ | 0.60071 |
| x_0 | 3.8750 |
| f | 6.5812 |
| $4.893 [\bar{\gamma}_f - \bar{\gamma}_\omega]$ | 47.39 |
| $4.893 [\bar{\gamma}_f + \bar{\gamma}_\omega]$ | 101.8 |

Hence the residue normalizations are given by $\bar{\gamma}_f = 15.25$ and $\bar{\gamma}_\omega = 5.55$.¹⁰

Thus as stated earlier we are left with one free parameter, namely d , which can be adjusted for a satisfactory description of the shrinkage-anti-shrinkage phenomenon of pp and $\bar{p}p$ elastic scattering. We find that a reasonable fit to the data is obtained with $d = 0.5 \text{ (GeV/c)}^{-2}$.

The comparison of this model with all available data for b , the logarithmic derivative of $d\sigma/dt$, is shown in Figs. 1a) and b). The fits tend to lie systematically slightly higher than the Serpukhov¹¹ and NAL¹² data for b in pp scattering. However, the data has a ± 0.3 systematic error which is not shown in the figures. The overall energy dependence of b in pp scattering is reproduced satisfactorily by the addition of ω and f^0 contributions to our previously parameterized pomeron amplitude. The trend of the data suggests that in our original⁶ choice of input data to determine

the parameters of the pomeron contribution, we may have chosen too large a value of b ($=11.4$) in the range $0.14 \leq |t| \leq 0.24(\text{GeV}/c)^2$. However, this does not affect our prediction for the energy dependence of the ratio of elastic to total cross section in a significant way. In Fig. 1b) it can be seen that the magnitude of the experimental values¹³ of $b_{\overline{pp}}$ and the slow anti-shrinkage is again well reproduced by the model.

We have calculated the ratio of the real to imaginary parts of the elastic amplitudes for pp and \overline{pp} scattering at various fixed t as a function of s . These predictions are given in Figs. 2a) and b). The curves for $t = 0$ are identical to those we obtained earlier.⁷ The recent results of the USA-USSR¹⁴ collaboration for pp scattering at $t = 0$ are shown on Fig. 2a). In addition, the previous data from the CERN-Rome group¹⁵ at the ISR are also shown. There is good agreement between the curve for $t = 0$ and the data. The other curves are new predictions of our model for the scattering amplitude away from the forward direction.¹⁶ Experimental data on both the pp and \overline{pp} cases will be of considerable interest.

We may conclude that the simplest Regge exchange contributions namely the ω and f^0 exchanges in addition to the pomeron amplitude are sufficient to account for the pre-asymptotic (20 to about 200 GeV) behavior of proton-proton and antiproton-proton scattering. The successful description of the energy and momentum dependence of these data at less than 200 GeV provides confidence in the accuracy of the Regge background contribution to the predicted energy dependence of $\sigma_{el}(s)$.

Figure 3a) shows the integrated elastic cross section $\sigma_{el}(s)$ for both pp and $\bar{p}p$ scattering versus s. Data from Serpukhov,¹⁷ NAL,¹⁸ and the ISR¹⁹ can be seen to follow closely the general trend of the pp curve. The curve lies a few percent below the data because of our original choice of too large a slope parameter b as mentioned earlier. Finally, Fig. 3b) shows the predicted ratio R defined to be $\sigma_{el}(s)/\sigma_{total}(s)$ plotted versus s. It can be seen that R(pp) and R($\bar{p}p$) both fall at low s, reach a minimum at s $\approx 300 \text{ GeV}^2$ and then tend to rise at high s. In the range accessible to the ISR, $500 \leq s \leq 3800$, there is a 7% increase in R. The accuracy²⁰ of the available data is not quite sufficient to observe the predicted increase in R.

In conclusion, we would like to emphasize that in Regge theories with cuts, such as those of Gribov,²¹ Frautschi and Margolis,²² and Dean,²³ it is very difficult to see how $\sigma_{el}(s)$ can increase with increasing energy. On the other hand, the Impact Picture¹ predicts not only that $\sigma_{el}(s)$ rises with energy, but also that it increases faster than σ_{total} . Therefore, the 7% rise in the ratio of elastic to total cross sections, if confirmed by experiment at the ISR, will represent a significant step forward in establishing the validity of the Impact Picture of high energy scattering. It is strongly urged that more accurate measurements be made in the near future to confront this prediction.

We would like to acknowledge stimulating discussions with Professors B. Kayser and G. Kane.

REFERENCES

*Work supported in part by the National Science Foundation under Grant No. GP-29463.

†Work supported by the U. S. Atomic Energy Commission under Contract No. AT(49-8)-3000.

‡Work supported in part by the U. S. Atomic Energy Commission under Contract No. AT(11-1)-3227.

¹H. Cheng and T. T. Wu, Phys. Rev. Letters 24, 1456 (1970); Phys. Letters 34B, 647 (1971).

²H. Cheng, J. K. Walker, and T. T. Wu, contribution to the XVI International Conference on High Energy Physics (1972).

³U. Amaldi et al., Phys. Letters 44B, 112 (1973).

⁴S. R. Amendolia et al., Phys. Letters 44B, 119 (1973).

⁵Aachen-CERN-Harvard-Geneva-Torino Collaboration presented by C. Rubbia at the XVI International Conference on High Energy Physics (1972).

⁶H. Cheng, J. K. Walker, and T. T. Wu, Phys. Letters 44B, 97 (1973).

⁷H. Cheng, J. K. Walker, and T. T. Wu, Phys. Letters 44B, 283 (1973).

⁸H. Cheng, J. K. Walker, and T. T. Wu, Phys. Rev. (to be published in Physical Review).

- ⁹V. Barger, K. Geer, and F. Halzen, CERN Report TH.1471 (1972).
- ¹⁰The ratio of the residue functions $\gamma_f(o)/\gamma_\omega(o) = 2.75$. This value has to be compared with unity which would be expected on the basis of exact exchange degeneracy. On the other hand the total pp cross section data clearly indicate that exchange degeneracy is not exact. It may be remarked that our other fit with $c' = 1$ allows a much milder (the above ratio is about 1.3) breaking of exchange degeneracy. We regard this as weak but positive evidence in favor of that choice for c' .
- ¹¹G. Beznogikh et al., Phys. Letters 30B (1969). The ISR data is from M. Holder et al., Phys. Letters 36B, 400 (1971); U. Amaldi et al., Phys. Letters 36B, 504 (1971); M. Holder et al., Phys. Letters 35B, 355 (1971).
- ¹²V. Bartenev et al., Phys. Rev. Letters 31, 1088 (1973).
- ¹³Yu. Antipov et al., contribution to the XVI International Conference on High Energy Physics (1972).
- ¹⁴V. Bartenev et al., Preprint NAL-Pub 73/54-EXP, 7100.036. Submitted to Phys. Rev. Letters for publication.
- ¹⁵U. Amaldi et al., Phys. Letters B43, 231 (1973) and CERN Report NP 73-19.
- ¹⁶We do not regard these predictions on the real to imaginary parts of the elastic amplitude away from the forward direction to be as reliable as our prediction for the rise of $R = \sigma_{el}(s)/\sigma_{total}(s)$. The increase in R emerges in a natural way from the Impact Picture whereas the real to imaginary ratios are very sensitive to the Regge background terms. However we believe that the predictions of the real to imaginary ratios are interesting and hopefully will stimulate their measurement.

- ¹⁷G. C. Beznogikh et al., contribution to the XVI International Conference on High Energy Physics (1972).
- ¹⁸J. W. Chapman et al., Phys. Rev. Letters 29, 1686 (1972); J. Dao et al., Phys. Rev. Letters 29, 1627 (1972); G. Charlton et al., contribution to the XVI International Conference on High Energy Physics (1972).
- ¹⁹See References 3 and 15.
- ²⁰In a measurement of $R = \sigma_{el}(s)/\sigma_{total}(s)$, errors in the luminosity need not enter if the measurements are made simultaneously at the one interaction region. In this way, it is hoped that the required accuracy may be achieved.
- ²¹V. N. Gribov, Zh. Eksp. Teor. Fiz. 53, 654 (1967) [JETP 26, 414 (1968)].
- ²²S. C. Frautschi and B. Margolis, Nuovo Cimento 56A, 1155 (1968).
- ²³N. W. Dean, Phys. Rev. 182, 1695 (1969).

FIGURE CAPTIONS

- Fig. 1a), b) The slope parameter, b , for proton-proton and anti-proton-proton elastic scattering is plotted versus the square of the center of mass energy s . The curves are for the two intervals of $|t|$ indicated. The average value of $|t|$ for each set of data is shown. Data from Serpukhov, NAL, and the ISR are shown as squares, crosses, and circles respectively.
- Fig. 2a), b) The predicted ratio of the real to imaginary parts of the proton-proton and antiproton-proton elastic scattering amplitudes are shown versus the square of the center of mass energy s . The data shown is for $|t| = 0 \text{ (GeV/c)}^2$.
- Fig. 3a) The predicted elastic cross section for proton-proton and antiproton-proton scattering is plotted versus the square of the center of mass energy $s \text{ (GeV)}^2$.
- Fig. 3b) The predicted ratio of the elastic to total cross section for proton-proton and antiproton-proton interactions is plotted versus the square of the center of mass energy $s \text{ (GeV)}^2$.





Fig. 1





Fig. 2





Fig. 3